

Quantum conditional relative entropy and quasi-factorization of the relative entropy

Ángela Capel (ICMAT-UAM)

Joint work with Angelo Lucia (U. Copenhagen)
and David Pérez-García (U. Complutense de Madrid).

Based on arXiv: 1705.03521 and 1804.09525



TQC2018, Sydney, 17 July 2018

CONTENTS

1 MOTIVATION

- QUANTUM DISSIPATIVE SYSTEMS
- CLASSICAL CASE

2 CONDITIONAL RELATIVE ENTROPY

- CONDITIONAL RELATIVE ENTROPY
- QUASI-FACTORIZATION FOR THE CONDITIONAL RELATIVE ENTROPY

3 QUANTUM SPIN LATTICES

- LOG-SOBOLEV CONSTANT

MOTIVATION




Figure: An open quantum many-body system.

- Interesting for information processing \Rightarrow Open (unavoidable interactions).
- Dynamics of S is dissipative!
- The continuous-time evolution of a state on S is given by a **quantum Markov semigroup**.

MOTIVATION




Figure: A quantum spin lattice system.

- Lattice $\Lambda \subset \mathbb{Z}^d$.
- To every site $x \in \Lambda$ we associate \mathcal{H}_x ($= \mathbb{C}^D$).
- The global Hilbert space associated to Λ is $\mathcal{H}_\Lambda = \bigotimes_{x \in \Lambda} \mathcal{H}_x$.

DISSIPATIVE QUANTUM SYSTEMS

DISSIPATIVE QUANTUM SYSTEMS

A **dissipative quantum system** is a 1-parameter continuous semigroup $\{\mathcal{T}_t^*\}_{t \geq 0}$ of completely positive, trace preserving (CPTP) maps (a.k.a. quantum channels) in \mathcal{S}_Λ .

$$\rho_\Lambda \xrightarrow{t} \rho_t := \mathcal{T}_t^*(\rho_\Lambda) = e^{t\mathcal{L}_\Lambda^*}(\rho_\Lambda) \xrightarrow{t \rightarrow \infty} \sigma_\Lambda$$

DISSIPATIVE QUANTUM SYSTEMS

DISSIPATIVE QUANTUM SYSTEMS

A **dissipative quantum system** is a 1-parameter continuous semigroup $\{\mathcal{T}_t^*\}_{t \geq 0}$ of completely positive, trace preserving (CPTP) maps (a.k.a. quantum channels) in \mathcal{S}_Λ .

$$\rho_\Lambda \xrightarrow{t} \rho_t := \mathcal{T}_t^*(\rho_\Lambda) = e^{t\mathcal{L}_\Lambda^*}(\rho_\Lambda) \xrightarrow{t \rightarrow \infty} \sigma_\Lambda$$

RAPID MIXING

We say that \mathcal{L}_Λ^* satisfies **rapid mixing** if

$$\sup_{\rho_\Lambda \in \mathcal{S}_\Lambda} \|\rho_t - \sigma_\Lambda\|_1 \leq \log(|\Lambda|) e^{-\gamma t}.$$

DISSIPATIVE QUANTUM SYSTEMS

DISSIPATIVE QUANTUM SYSTEMS

A **dissipative quantum system** is a 1-parameter continuous semigroup $\{\mathcal{T}_t^*\}_{t \geq 0}$ of completely positive, trace preserving (CPTP) maps (a.k.a. quantum channels) in \mathcal{S}_Λ .

$$\rho_\Lambda \xrightarrow{t} \rho_t := \mathcal{T}_t^*(\rho_\Lambda) = e^{t\mathcal{L}_\Lambda^*}(\rho_\Lambda) \xrightarrow{t \rightarrow \infty} \sigma_\Lambda$$

RAPID MIXING

We say that \mathcal{L}_Λ^* satisfies **rapid mixing** if

$$\sup_{\rho_\Lambda \in \mathcal{S}_\Lambda} \|\rho_t - \sigma_\Lambda\|_1 \leq \log(|\Lambda|)e^{-\gamma t}.$$

PROBLEM

Find examples of rapid mixing!

DISSIPATIVE QUANTUM SYSTEMS

DISSIPATIVE QUANTUM SYSTEMS

A **dissipative quantum system** is a 1-parameter continuous semigroup $\{\mathcal{T}_t^*\}_{t \geq 0}$ of completely positive, trace preserving (CPTP) maps (a.k.a. quantum channels) in \mathcal{S}_Λ .

$$\rho_\Lambda \xrightarrow{t} \rho_t := \mathcal{T}_t^*(\rho_\Lambda) = e^{t\mathcal{L}_\Lambda^*}(\rho_\Lambda) \xrightarrow{t \rightarrow \infty} \sigma_\Lambda$$

RAPID MIXING

We say that \mathcal{L}_Λ^* satisfies **rapid mixing** if

$$\sup_{\rho_\Lambda \in \mathcal{S}_\Lambda} \|\rho_t - \sigma_\Lambda\|_1 \leq \log(|\Lambda|)e^{-\gamma t}.$$

PROBLEM

Find examples of rapid mixing!

LOG-SOBOLEV CONSTANT

LOG-SOBOLEV CONSTANT

Let $\mathcal{L}_\Lambda^* : \mathcal{S}_\Lambda \rightarrow \mathcal{S}_\Lambda$ be a primitive reversible Lindbladian with stationary state σ_Λ . We define the **log-Sobolev constant** of \mathcal{L}_Λ^* by

$$\alpha(\mathcal{L}_\Lambda^*) := \inf_{\rho_\Lambda \in \mathcal{S}_\Lambda} \frac{-\operatorname{tr}[\mathcal{L}_\Lambda^*(\rho_\Lambda)(\log \rho_\Lambda - \log \sigma_\Lambda)]}{2D(\rho_\Lambda || \sigma_\Lambda)}$$

If $\alpha(\mathcal{L}_\Lambda^*) > 0$:

$$D(\rho_t || \sigma_\Lambda) \leq D(\rho_\Lambda || \sigma_\Lambda) e^{-2\alpha(\mathcal{L}_\Lambda^*)t},$$

LOG-SOBOLEV CONSTANT

LOG-SOBOLEV CONSTANT

Let $\mathcal{L}_\Lambda^* : \mathcal{S}_\Lambda \rightarrow \mathcal{S}_\Lambda$ be a primitive reversible Lindbladian with stationary state σ_Λ . We define the **log-Sobolev constant** of \mathcal{L}_Λ^* by

$$\alpha(\mathcal{L}_\Lambda^*) := \inf_{\rho_\Lambda \in \mathcal{S}_\Lambda} \frac{-\text{tr}[\mathcal{L}_\Lambda^*(\rho_\Lambda)(\log \rho_\Lambda - \log \sigma_\Lambda)]}{2D(\rho_\Lambda || \sigma_\Lambda)}$$

If $\alpha(\mathcal{L}_\Lambda^*) > 0$:

$$D(\rho_t || \sigma_\Lambda) \leq D(\rho_\Lambda || \sigma_\Lambda) e^{-2\alpha(\mathcal{L}_\Lambda^*)t},$$

and putting this together with **Pinsker's inequality**, we have:

$$\|\rho_t - \sigma_\Lambda\|_1 \leq \sqrt{2D(\rho_t || \sigma_\Lambda)} e^{-\alpha(\mathcal{L}_\Lambda^*)t} \leq \sqrt{2\log(1/\sigma_{\min})} e^{-\alpha(\mathcal{L}_\Lambda^*)t}.$$

LOG-SOBOLEV CONSTANT

LOG-SOBOLEV CONSTANT

Let $\mathcal{L}_\Lambda^* : \mathcal{S}_\Lambda \rightarrow \mathcal{S}_\Lambda$ be a primitive reversible Lindbladian with stationary state σ_Λ . We define the **log-Sobolev constant** of \mathcal{L}_Λ^* by

$$\alpha(\mathcal{L}_\Lambda^*) := \inf_{\rho_\Lambda \in \mathcal{S}_\Lambda} \frac{-\text{tr}[\mathcal{L}_\Lambda^*(\rho_\Lambda)(\log \rho_\Lambda - \log \sigma_\Lambda)]}{2D(\rho_\Lambda || \sigma_\Lambda)}$$

If $\alpha(\mathcal{L}_\Lambda^*) > 0$:

$$D(\rho_t || \sigma_\Lambda) \leq D(\rho_\Lambda || \sigma_\Lambda) e^{-2\alpha(\mathcal{L}_\Lambda^*)t},$$

and putting this together with **Pinsker's inequality**, we have:

$$\|\rho_t - \sigma_\Lambda\|_1 \leq \sqrt{2D(\rho_\Lambda || \sigma_\Lambda)} e^{-\alpha(\mathcal{L}_\Lambda^*)t} \leq \sqrt{2 \log(1/\sigma_{\min})} e^{-\alpha(\mathcal{L}_\Lambda^*)t}.$$

Log-Sobolev constant \Rightarrow Rapid mixing.

LOG-SOBOLEV CONSTANT

LOG-SOBOLEV CONSTANT

Let $\mathcal{L}_\Lambda^* : \mathcal{S}_\Lambda \rightarrow \mathcal{S}_\Lambda$ be a primitive reversible Lindbladian with stationary state σ_Λ . We define the **log-Sobolev constant** of \mathcal{L}_Λ^* by

$$\alpha(\mathcal{L}_\Lambda^*) := \inf_{\rho_\Lambda \in \mathcal{S}_\Lambda} \frac{-\text{tr}[\mathcal{L}_\Lambda^*(\rho_\Lambda)(\log \rho_\Lambda - \log \sigma_\Lambda)]}{2D(\rho_\Lambda || \sigma_\Lambda)}$$

If $\alpha(\mathcal{L}_\Lambda^*) > 0$:

$$D(\rho_t || \sigma_\Lambda) \leq D(\rho_\Lambda || \sigma_\Lambda) e^{-2\alpha(\mathcal{L}_\Lambda^*)t},$$

and putting this together with **Pinsker's inequality**, we have:

$$\|\rho_t - \sigma_\Lambda\|_1 \leq \sqrt{2D(\rho_t || \sigma_\Lambda)} e^{-\alpha(\mathcal{L}_\Lambda^*)t} \leq \sqrt{2\log(1/\sigma_{\min})} e^{-\alpha(\mathcal{L}_\Lambda^*)t}.$$

Log-Sobolev constant \Rightarrow Rapid mixing.

PROBLEM

Find positive log-Sobolev constants!

LOG-SOBOLEV CONSTANT

LOG-SOBOLEV CONSTANT

Let $\mathcal{L}_\Lambda^* : \mathcal{S}_\Lambda \rightarrow \mathcal{S}_\Lambda$ be a primitive reversible Lindbladian with stationary state σ_Λ . We define the **log-Sobolev constant** of \mathcal{L}_Λ^* by

$$\alpha(\mathcal{L}_\Lambda^*) := \inf_{\rho_\Lambda \in \mathcal{S}_\Lambda} \frac{-\text{tr}[\mathcal{L}_\Lambda^*(\rho_\Lambda)(\log \rho_\Lambda - \log \sigma_\Lambda)]}{2D(\rho_\Lambda || \sigma_\Lambda)}$$

If $\alpha(\mathcal{L}_\Lambda^*) > 0$:

$$D(\rho_t || \sigma_\Lambda) \leq D(\rho_\Lambda || \sigma_\Lambda) e^{-2\alpha(\mathcal{L}_\Lambda^*)t},$$

and putting this together with **Pinsker's inequality**, we have:

$$\|\rho_t - \sigma_\Lambda\|_1 \leq \sqrt{2D(\rho_t || \sigma_\Lambda)} e^{-\alpha(\mathcal{L}_\Lambda^*)t} \leq \sqrt{2\log(1/\sigma_{\min})} e^{-\alpha(\mathcal{L}_\Lambda^*)t}.$$

Log-Sobolev constant \Rightarrow Rapid mixing.

PROBLEM

Find positive log-Sobolev constants!

CLASSICAL CASE

CLASSICAL ENTROPY AND CONDITIONAL ENTROPY

Consider a probability space $(\Omega, \mathcal{F}, \mu)$ and define, for every $f > 0$, the **entropy** of f by

$$\text{Ent}_\mu(f) = \mu(f \log f) - \mu(f) \log \mu(f).$$

Given a σ -algebra $\mathcal{G} \subseteq \mathcal{F}$, we define the **conditional entropy** of f in \mathcal{G} by

$$\text{Ent}_\mu(f | \mathcal{G}) = \mu(f \log f | \mathcal{G}) - \mu(f | \mathcal{G}) \log \mu(f | \mathcal{G}).$$

LEMMA, Dai Pra et al. '02

Let $(\Omega, \mathcal{F}, \mu)$ be a probability space, and $\mathcal{F}_1, \mathcal{F}_2$ sub- σ -algebras of \mathcal{F} . Suppose that there exists a probability measure $\bar{\mu}$ that makes \mathcal{F}_1 and \mathcal{F}_2 independent, $\mu \ll \bar{\mu}$ and $\mu | \mathcal{F}_i = \bar{\mu} | \mathcal{F}_i$ for $i = 1, 2$. Then, for every $f \geq 0$ such that $f \log f \in L^1(\mu)$ and $\mu(f) = 1$,

$$\text{Ent}_\mu(f) \leq \frac{1}{1 - 4\|h - 1\|_\infty} \mu[\text{Ent}_\mu(f | \mathcal{F}_1) + \text{Ent}_\mu(f | \mathcal{F}_2)],$$

where $h = \frac{d\mu}{d\bar{\mu}}$.

CLASSICAL CASE

CLASSICAL ENTROPY AND CONDITIONAL ENTROPY

Consider a probability space $(\Omega, \mathcal{F}, \mu)$ and define, for every $f > 0$, the **entropy** of f by

$$\text{Ent}_\mu(f) = \mu(f \log f) - \mu(f) \log \mu(f).$$

Given a σ -algebra $\mathcal{G} \subseteq \mathcal{F}$, we define the **conditional entropy** of f in \mathcal{G} by

$$\text{Ent}_\mu(f | \mathcal{G}) = \mu(f \log f | \mathcal{G}) - \mu(f | \mathcal{G}) \log \mu(f | \mathcal{G}).$$

LEMMA, Dai Pra et al. '02

Let $(\Omega, \mathcal{F}, \mu)$ be a probability space, and $\mathcal{F}_1, \mathcal{F}_2$ sub- σ -algebras of \mathcal{F} . Suppose that there exists a probability measure $\bar{\mu}$ that makes \mathcal{F}_1 and \mathcal{F}_2 independent, $\mu \ll \bar{\mu}$ and $\mu | \mathcal{F}_i = \bar{\mu} | \mathcal{F}_i$ for $i = 1, 2$. Then, for every $f \geq 0$ such that $f \log f \in L^1(\mu)$ and $\mu(f) = 1$,

$$\text{Ent}_\mu(f) \leq \frac{1}{1 - 4\|h - 1\|_\infty} \mu[\text{Ent}_\mu(f | \mathcal{F}_1) + \text{Ent}_\mu(f | \mathcal{F}_2)],$$

where $h = \frac{d\mu}{d\bar{\mu}}$.

CLASSICAL CASE

CLASSICAL CASE

$$\text{Ent}_\mu(f) \leq \frac{1}{1 - 4\|h - 1\|_\infty} \mu [\text{Ent}_\mu(f | \mathcal{F}_1) + \text{Ent}_\mu(f | \mathcal{F}_2)],$$

where $h = \frac{d\mu}{d\bar{\mu}}$.

PROBLEM

Let $\mathcal{H}_{ABC} = \mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$ and $\rho_{ABC}, \sigma_{ABC} \in S_{ABC}$. Can we prove something like

$$D(\rho_{ABC} || \sigma_{ABC}) \leq \xi(\sigma_{ABC}) [D_{AB}(\rho_{ABC} || \sigma_{ABC}) + D_{BC}(\rho_{ABC} || \sigma_{ABC})] ?$$

Yes! (We will see how later)

CLASSICAL CASE

CLASSICAL CASE

$$\text{Ent}_\mu(f) \leq \frac{1}{1 - 4\|h - 1\|_\infty} \mu [\text{Ent}_\mu(f | \mathcal{F}_1) + \text{Ent}_\mu(f | \mathcal{F}_2)],$$

where $h = \frac{d\mu}{d\bar{\mu}}$.

PROBLEM

Let $\mathcal{H}_{ABC} = \mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$ and $\rho_{ABC}, \sigma_{ABC} \in S_{ABC}$. Can we prove something like

$$D(\rho_{ABC} || \sigma_{ABC}) \leq \xi(\sigma_{ABC}) [D_{AB}(\rho_{ABC} || \sigma_{ABC}) + D_{BC}(\rho_{ABC} || \sigma_{ABC})] ?$$

Yes! (We will see how later)

RELATIVE ENTROPY

QUANTUM RELATIVE ENTROPY

Let $\rho_\Lambda, \sigma_\Lambda \in \mathcal{S}_\Lambda$. The **quantum relative entropy** of ρ_Λ and σ_Λ is defined by:

$$D(\rho_\Lambda || \sigma_\Lambda) = \text{tr} [\rho_\Lambda (\log \rho_\Lambda - \log \sigma_\Lambda)].$$

PROPERTIES OF THE RELATIVE ENTROPY

Let $\mathcal{H}_{AB} = \mathcal{H}_A \otimes \mathcal{H}_B$ and $\rho_{AB}, \sigma_{AB} \in \mathcal{S}_{AB}$. The following properties hold:

- 1 **Continuity.** $\rho_{AB} \mapsto D(\rho_{AB} || \sigma_{AB})$ is continuous.
- 2 **Additivity.** $D(\rho_A \otimes \rho_B || \sigma_A \otimes \sigma_B) = D(\rho_A || \sigma_A) + D(\rho_B || \sigma_B)$.
- 3 **Superadditivity.** $D(\rho_{AB} || \sigma_A \otimes \sigma_B) \geq D(\rho_A || \sigma_A) + D(\rho_B || \sigma_B)$.
- 4 **Monotonicity.** $D(\rho_{AB} || \sigma_{AB}) \geq D(T(\rho_{AB}) || T(\sigma_{AB}))$ for every quantum channel T .

CHARACTERIZATION OF THE RELATIVE ENTROPY, Wilming et al. '17

If $f : \mathcal{S}_{AB} \times \mathcal{S}_{AB} \rightarrow \mathbb{R}_0^+$ satisfies 1 – 4, then f is the relative entropy.

RELATIVE ENTROPY

QUANTUM RELATIVE ENTROPY

Let $\rho_\Lambda, \sigma_\Lambda \in \mathcal{S}_\Lambda$. The **quantum relative entropy** of ρ_Λ and σ_Λ is defined by:

$$D(\rho_\Lambda || \sigma_\Lambda) = \text{tr} [\rho_\Lambda (\log \rho_\Lambda - \log \sigma_\Lambda)].$$

PROPERTIES OF THE RELATIVE ENTROPY

Let $\mathcal{H}_{AB} = \mathcal{H}_A \otimes \mathcal{H}_B$ and $\rho_{AB}, \sigma_{AB} \in \mathcal{S}_{AB}$. The following properties hold:

- 1 **Continuity.** $\rho_{AB} \mapsto D(\rho_{AB} || \sigma_{AB})$ is continuous.
- 2 **Additivity.** $D(\rho_A \otimes \rho_B || \sigma_A \otimes \sigma_B) = D(\rho_A || \sigma_A) + D(\rho_B || \sigma_B)$.
- 3 **Superadditivity.** $D(\rho_{AB} || \sigma_A \otimes \sigma_B) \geq D(\rho_A || \sigma_A) + D(\rho_B || \sigma_B)$.
- 4 **Monotonicity.** $D(\rho_{AB} || \sigma_{AB}) \geq D(T(\rho_{AB}) || T(\sigma_{AB}))$ for every quantum channel T .

CHARACTERIZATION OF THE RELATIVE ENTROPY, Wilming et al. '17

If $f : \mathcal{S}_{AB} \times \mathcal{S}_{AB} \rightarrow \mathbb{R}_0^+$ satisfies 1 – 4, then f is the relative entropy.

CONDITIONAL RELATIVE ENTROPY

CONDITIONAL RELATIVE ENTROPY

Let $\mathcal{H}_{AB} = \mathcal{H}_A \otimes \mathcal{H}_B$. We define a **conditional relative entropy** in A as a function

$$D_A(\cdot || \cdot) : \mathcal{S}_{AB} \times \mathcal{S}_{AB} \rightarrow \mathbb{R}_0^+$$

verifying the following properties for every $\rho_{AB}, \sigma_{AB} \in \mathcal{S}_{AB}$:

① **Continuity:** The map $\rho_{AB} \mapsto D_A(\rho_{AB} || \sigma_{AB})$ is continuous.

② **Non-negativity:** $D_A(\rho_{AB} || \sigma_{AB}) \geq 0$ and

$$(2.1) \quad D_A(\rho_{AB} || \sigma_{AB}) = 0 \text{ if, and only if, } \rho_{AB} = \sigma_{AB}^{1/2} \sigma_B^{-1/2} \rho_{AB} \sigma_B^{-1/2} \sigma_{AB}^{1/2}.$$

③ **Semi-superadditivity:** $D_A(\rho_{AB} || \sigma_A \otimes \sigma_B) \geq D(\rho_A || \sigma_A)$ and

$$(3.1) \quad \text{Semi-additivity: if } \rho_{AB} = \rho_A \otimes \rho_B, \\ D_A(\rho_A \otimes \rho_B || \sigma_A \otimes \sigma_B) = D(\rho_A || \sigma_A).$$

④ **Semi-motonicity:** For every quantum channel \mathcal{T} ,

$$D_A(\mathcal{T}(\rho_{AB}) || \mathcal{T}(\sigma_{AB})) + D_B((\text{tr}_A \circ \mathcal{T})(\rho_{AB}) || (\text{tr}_A \circ \mathcal{T})(\sigma_{AB})) \\ \leq D_A(\rho_{AB} || \sigma_{AB}) + D_B(\text{tr}_A(\rho_{AB}) || \text{tr}_A(\sigma_{AB})).$$

REMARK

Consider for every $\rho_{AB}, \sigma_{AB} \in \mathcal{S}_{AB}$

$$D_{A,B}^+(\rho_{AB}||\sigma_{AB}) = D_A(\rho_{AB}||\sigma_{AB}) + D_B(\rho_{AB}||\sigma_{AB}).$$

Then, $D_{A,B}^+$ verifies the following properties:

- 1 **Continuity:** $\rho_{AB} \mapsto D_{A,B}^+(\rho_{AB}||\sigma_{AB})$ is continuous.
- 2 **Additivity:** $D_{A,B}^+(\rho_A \otimes \rho_B || \sigma_A \otimes \sigma_B) = D(\rho_A || \sigma_A) + D(\rho_B || \sigma_B)$.
- 3 **Superadditivity:** $D_{A,B}^+(\rho_{AB} || \sigma_A \otimes \sigma_B) \geq D(\rho_A || \sigma_A) + D(\rho_B || \sigma_B)$.

However, it does not satisfy the property of monotonicity.

AXIOMATIC CHARACTERIZATION OF THE CONDITIONAL RELATIVE ENTROPY

The only possible conditional relative entropy is given by:

$$D_A(\rho_{AB}||\sigma_{AB}) = D(\rho_{AB}||\sigma_{AB}) - D(\rho_B||\sigma_B)$$

for every $\rho_{AB}, \sigma_{AB} \in \mathcal{S}_{AB}$.

REMARK

Consider for every $\rho_{AB}, \sigma_{AB} \in \mathcal{S}_{AB}$

$$D_{A,B}^+(\rho_{AB}||\sigma_{AB}) = D_A(\rho_{AB}||\sigma_{AB}) + D_B(\rho_{AB}||\sigma_{AB}).$$

Then, $D_{A,B}^+$ verifies the following properties:

- 1 **Continuity:** $\rho_{AB} \mapsto D_{A,B}^+(\rho_{AB}||\sigma_{AB})$ is continuous.
- 2 **Additivity:** $D_{A,B}^+(\rho_A \otimes \rho_B || \sigma_A \otimes \sigma_B) = D(\rho_A || \sigma_A) + D(\rho_B || \sigma_B)$.
- 3 **Superadditivity:** $D_{A,B}^+(\rho_{AB} || \sigma_A \otimes \sigma_B) \geq D(\rho_A || \sigma_A) + D(\rho_B || \sigma_B)$.

However, it does not satisfy the property of monotonicity.

AXIOMATIC CHARACTERIZATION OF THE CONDITIONAL RELATIVE ENTROPY

The only possible conditional relative entropy is given by:

$$D_A(\rho_{AB}||\sigma_{AB}) = D(\rho_{AB}||\sigma_{AB}) - D(\rho_B||\sigma_B)$$

for every $\rho_{AB}, \sigma_{AB} \in \mathcal{S}_{AB}$.




Figure: Choice of indices in $\mathcal{H}_{ABC} = \mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$.

Result of **quasi-factorization** of the relative entropy, for every $\rho_{ABC}, \sigma_{ABC} \in \mathcal{S}_{ABC}$:

$$D(\rho_{ABC} || \sigma_{ABC}) \leq \xi(\sigma_{ABC}) [D_{AB}(\rho_{ABC} || \sigma_{ABC}) + D_{BC}(\rho_{ABC} || \sigma_{ABC})],$$

where $\xi(\sigma_{ABC})$ depends only on σ_{ABC} and measures how far σ_{AC} is from $\sigma_A \otimes \sigma_C$.

QUASI-FACTORIZATION FOR THE CRE

Let $\mathcal{H}_{ABC} = \mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$ and $\rho_{ABC}, \sigma_{ABC} \in \mathcal{S}_{ABC}$. Then, the following inequality holds

$$D(\rho_{ABC} \parallel \sigma_{ABC}) \leq \frac{1}{1 - 2\|H(\sigma_{AC})\|_\infty} [D_{AB}(\rho_{ABC} \parallel \sigma_{ABC}) + D_{BC}(\rho_{ABC} \parallel \sigma_{ABC})],$$

where

$$H(\sigma_{AC}) = \sigma_A^{-1/2} \otimes \sigma_C^{-1/2} \sigma_{AC} \sigma_A^{-1/2} \otimes \sigma_C^{-1/2} - \mathbb{1}_{AC}.$$

Note that $H(\sigma_{AC}) = 0$ if σ_{AC} is a tensor product between A and C .

CLASSICAL CASE

$$\text{Ent}_\mu(f) \leq \frac{1}{1 - 4\|h - 1\|_\infty} \mu [\text{Ent}_\mu(f \mid \mathcal{F}_1) + \text{Ent}_\mu(f \mid \mathcal{F}_2)],$$

where $h = \frac{d\mu}{d\bar{\mu}}$.

QUASI-FACTORIZATION FOR THE CRE

Let $\mathcal{H}_{ABC} = \mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$ and $\rho_{ABC}, \sigma_{ABC} \in \mathcal{S}_{ABC}$. Then, the following inequality holds

$$D(\rho_{ABC} || \sigma_{ABC}) \leq \frac{1}{1 - 2\|H(\sigma_{AC})\|_\infty} [D_{AB}(\rho_{ABC} || \sigma_{ABC}) + D_{BC}(\rho_{ABC} || \sigma_{ABC})],$$

where

$$H(\sigma_{AC}) = \sigma_A^{-1/2} \otimes \sigma_C^{-1/2} \sigma_{AC} \sigma_A^{-1/2} \otimes \sigma_C^{-1/2} - \mathbb{1}_{AC}.$$

Note that $H(\sigma_{AC}) = 0$ if σ_{AC} is a tensor product between A and C .

CLASSICAL CASE

$$\text{Ent}_\mu(f) \leq \frac{1}{1 - 4\|h - 1\|_\infty} \mu [\text{Ent}_\mu(f | \mathcal{F}_1) + \text{Ent}_\mu(f | \mathcal{F}_2)],$$

where $h = \frac{d\mu}{d\bar{\mu}}$.

This result is equivalent to:

$$\boxed{(1 + 2\|H(\sigma_{AB})\|_\infty)D(\rho_{AB}||\sigma_{AB}) \geq D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B)}.$$

Recall:

- **Superadditivity.** $D(\rho_{AB}||\sigma_A \otimes \sigma_B) \geq D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B)$.

This result is equivalent to:

$$\boxed{(1 + 2\|H(\sigma_{AB})\|_\infty)D(\rho_{AB}||\sigma_{AB}) \geq D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B)}.$$

Recall:

- **Superadditivity.** $D(\rho_{AB}||\sigma_A \otimes \sigma_B) \geq D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B)$.

Due to:

- **Monotonicity.** $D(\rho_{AB}||\sigma_{AB}) \geq D(T(\rho_{AB})||T(\sigma_{AB}))$ for every quantum channel T .

we have

$$2D(\rho_{AB}||\sigma_{AB}) \geq D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B).$$

This result is equivalent to:

$$\boxed{(1 + 2\|H(\sigma_{AB})\|_\infty)D(\rho_{AB}||\sigma_{AB}) \geq D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B)}.$$

Recall:

- **Superadditivity.** $D(\rho_{AB}||\sigma_A \otimes \sigma_B) \geq D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B)$.

Due to:

- **Monotonicity.** $D(\rho_{AB}||\sigma_{AB}) \geq D(T(\rho_{AB})||T(\sigma_{AB}))$ for every quantum channel T .

we have

$$2D(\rho_{AB}||\sigma_{AB}) \geq D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B).$$

RELATION WITH THE CLASSICAL CASE




Figure: Identification between classical and quantum quantities when the states considered are classical.

SKETCH OF THE PROOF OF QUASI-FACTORIZATION

$$(1 + 2\|H(\sigma_{AB})\|_\infty)D(\rho_{AB}||\sigma_{AB}) \geq D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B).$$

STEP 1

$$D(\rho_{AB}||\sigma_{AB}) \geq D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B) - \log \text{tr } M, \quad (1)$$

where $M = \exp[\log \sigma_{AB} - \log \sigma_A \otimes \sigma_B + \log \rho_A \otimes \rho_B]$.

SKETCH OF THE PROOF OF QUASI-FACTORIZATION

$$(1 + 2\|H(\sigma_{AB})\|_\infty)D(\rho_{AB}||\sigma_{AB}) \geq D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B).$$

STEP 1

$$D(\rho_{AB}||\sigma_{AB}) \geq D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B) - \log \operatorname{tr} M, \quad (1)$$

where $M = \exp[\log \sigma_{AB} - \log \sigma_A \otimes \sigma_B + \log \rho_A \otimes \rho_B]$.

STEP 2

$$\log \operatorname{tr} M \leq \operatorname{tr}[L(\sigma_{AB}) (\rho_A - \sigma_A) \otimes (\rho_B - \sigma_B)], \quad (2)$$

where

$$L(\sigma_{AB}) = \mathcal{T}_{\sigma_A \otimes \sigma_B}(\sigma_{AB}) - \mathbb{1}_{AB}.$$

SKETCH OF THE PROOF OF QUASI-FACTORIZATION

$$(1 + 2\|H(\sigma_{AB})\|_\infty)D(\rho_{AB}||\sigma_{AB}) \geq D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B).$$

STEP 1

$$D(\rho_{AB}||\sigma_{AB}) \geq D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B) - \log \text{tr } M, \quad (1)$$

where $M = \exp[\log \sigma_{AB} - \log \sigma_A \otimes \sigma_B + \log \rho_A \otimes \rho_B]$.

STEP 2

$$\log \text{tr } M \leq \text{tr}[L(\sigma_{AB}) (\rho_A - \sigma_A) \otimes (\rho_B - \sigma_B)], \quad (2)$$

where

$$L(\sigma_{AB}) = \mathcal{T}_{\sigma_A \otimes \sigma_B}(\sigma_{AB}) - \mathbb{1}_{AB}.$$

THEOREM (Lieb, '73)

Let g a positive operator, and define

$$\mathcal{T}_g(f) = \int_0^\infty dt (g+t)^{-1} f (g+t)^{-1}.$$

\mathcal{T}_g is positive-semidefinite if g is. We have that

$$\text{tr}[\exp(-f+g+h)] \leq \text{tr}[e^h \mathcal{T}_{ef}(e^g)].$$

LEMMA (Sutter et al., '17)

For $f \in \mathcal{S}_{AB}$ and $g \in \mathcal{A}_{AB}$ the following holds:

$$\mathcal{T}_g(f) = \int_{-\infty}^{\infty} dt \beta_0(t) g^{-\frac{1-it}{2}} f g^{-\frac{-1+it}{2}},$$

with

$$\beta_0(t) = \frac{\pi}{2} (\cosh(\pi t) + 1)^{-1}.$$

THEOREM (Lieb, '73)

Let g a positive operator, and define

$$\mathcal{T}_g(f) = \int_0^\infty dt (g+t)^{-1} f (g+t)^{-1}.$$

\mathcal{T}_g is positive-semidefinite if g is. We have that

$$\text{tr}[\exp(-f + g + h)] \leq \text{tr}\left[e^h \mathcal{T}_{e^f}(e^g)\right].$$

LEMMA (Sutter et al., '17)

For $f \in \mathcal{S}_{AB}$ and $g \in \mathcal{A}_{AB}$ the following holds:

$$\mathcal{T}_g(f) = \int_{-\infty}^{\infty} dt \beta_0(t) g^{\frac{-1-it}{2}} f g^{\frac{-1+it}{2}},$$

with

$$\beta_0(t) = \frac{\pi}{2} (\cosh(\pi t) + 1)^{-1}.$$

STEP 3

$$\mathrm{tr}[L(\sigma_{AB})(\rho_A - \sigma_A) \otimes (\rho_B - \sigma_B)] \leq 2\|L(\sigma_{AB})\|_\infty D(\rho_{AB}||\sigma_{AB}). \quad (3)$$

Hölder's inequality + Tensorization of Schatten norms + Pinsker's inequality + Data Processing inequality.

STEP 3

$$\mathrm{tr}[L(\sigma_{AB})(\rho_A - \sigma_A) \otimes (\rho_B - \sigma_B)] \leq 2\|L(\sigma_{AB})\|_\infty D(\rho_{AB}||\sigma_{AB}). \quad (3)$$

Hölder's inequality + Tensorization of Schatten norms + Pinsker's inequality + Data Processing inequality.

Step 4

$$\|L(\sigma_{AB})\|_\infty \leq \left\| \sigma_A^{-1/2} \otimes \sigma_B^{-1/2} \sigma_{AB} \sigma_A^{-1/2} \otimes \sigma_B^{-1/2} - \mathbb{1}_{AB} \right\|_\infty. \quad (4)$$

STEP 3

$$\mathrm{tr}[L(\sigma_{AB})(\rho_A - \sigma_A) \otimes (\rho_B - \sigma_B)] \leq 2\|L(\sigma_{AB})\|_\infty D(\rho_{AB}||\sigma_{AB}). \quad (3)$$

Hölder's inequality + Tensorization of Schatten norms + Pinsker's inequality + Data Processing inequality.

Step 4

$$\|L(\sigma_{AB})\|_\infty \leq \left\| \sigma_A^{-1/2} \otimes \sigma_B^{-1/2} \sigma_{AB} \sigma_A^{-1/2} \otimes \sigma_B^{-1/2} - \mathbb{1}_{AB} \right\|_\infty. \quad (4)$$

QUANTUM SPIN LATTICES




Figure: A quantum spin lattice system Λ and $A, B \subseteq \Lambda$ such that $A \cup B = \Lambda$.

PROBLEM

For a certain \mathcal{L}_Λ^* , can we prove $\alpha(\mathcal{L}_\Lambda^*) > 0$ using the result of quasi-factorization of the relative entropy?

GENERAL QUASI-FACTORIZATION FOR σ A TENSOR PRODUCT

Let $\mathcal{H}_\Lambda = \bigotimes_{x \in \Lambda} \mathcal{H}_x$ and $\rho_\Lambda, \sigma_\Lambda \in \mathcal{S}_\Lambda$ such that $\sigma_\Lambda = \bigotimes_{x \in \Lambda} \sigma_x$. The following inequality holds:

$$D(\rho_\Lambda || \sigma_\Lambda) \leq \sum_{x \in \Lambda} D_x(\rho_\Lambda || \sigma_\Lambda). \quad (5)$$

The **heat-bath dynamics**, with product fixed point, has a positive log-Sobolev constant.

GENERAL QUASI-FACTORIZATION FOR σ A TENSOR PRODUCT

Let $\mathcal{H}_\Lambda = \bigotimes_{x \in \Lambda} \mathcal{H}_x$ and $\rho_\Lambda, \sigma_\Lambda \in \mathcal{S}_\Lambda$ such that $\sigma_\Lambda = \bigotimes_{x \in \Lambda} \sigma_x$. The following inequality holds:

$$D(\rho_\Lambda || \sigma_\Lambda) \leq \sum_{x \in \Lambda} D_x(\rho_\Lambda || \sigma_\Lambda). \quad (5)$$

The **heat-bath dynamics**, with product fixed point, has a positive log-Sobolev constant.

Consider the local and global Lindbladians

$$\mathcal{L}_x^* := \mathbb{E}_x^* - \mathbb{1}_\Lambda, \quad \mathcal{L}_\Lambda^* = \sum_{x \in \Lambda} \mathcal{L}_x^*$$

Since

$$\mathbb{E}_x^*(\rho_\Lambda) = \sigma_\Lambda^{1/2} \sigma_{x^c}^{-1/2} \rho_{x^c} \sigma_{x^c}^{-1/2} \sigma_\Lambda^{1/2} = \sigma_x \otimes \rho_{x^c}$$

for every $\rho_\Lambda \in \mathcal{S}_\Lambda$, we have

$$\mathcal{L}_\Lambda^*(\rho_\Lambda) = \sum_{x \in \Lambda} (\sigma_x \otimes \rho_{x^c} - \rho_\Lambda).$$

GENERAL QUASI-FACTORIZATION FOR σ A TENSOR PRODUCT

Let $\mathcal{H}_\Lambda = \bigotimes_{x \in \Lambda} \mathcal{H}_x$ and $\rho_\Lambda, \sigma_\Lambda \in \mathcal{S}_\Lambda$ such that $\sigma_\Lambda = \bigotimes_{x \in \Lambda} \sigma_x$. The following inequality holds:

$$D(\rho_\Lambda || \sigma_\Lambda) \leq \sum_{x \in \Lambda} D_x(\rho_\Lambda || \sigma_\Lambda). \quad (5)$$

The **heat-bath dynamics**, with product fixed point, has a positive log-Sobolev constant.

Consider the local and global Lindbladians

$$\mathcal{L}_x^* := \mathbb{E}_x^* - \mathbb{1}_\Lambda, \quad \mathcal{L}_\Lambda^* = \sum_{x \in \Lambda} \mathcal{L}_x^*$$

Since

$$\mathbb{E}_x^*(\rho_\Lambda) = \sigma_\Lambda^{1/2} \sigma_{x^c}^{-1/2} \rho_{x^c} \sigma_{x^c}^{-1/2} \sigma_\Lambda^{1/2} = \sigma_x \otimes \rho_{x^c}$$

for every $\rho_\Lambda \in \mathcal{S}_\Lambda$, we have

$$\mathcal{L}_\Lambda^*(\rho_\Lambda) = \sum_{x \in \Lambda} (\sigma_x \otimes \rho_{x^c} - \rho_\Lambda).$$

LOG-SOBOLEV CONSTANT

CONDITIONAL LOG-SOBOLEV CONSTANT

For $x \in \Lambda$, we define the **conditional log-Sobolev constant** of \mathcal{L}_Λ^* in x by

$$\alpha_\Lambda(\mathcal{L}_x^*) := \inf_{\rho_\Lambda \in \mathcal{S}_\Lambda} \frac{-\operatorname{tr}[\mathcal{L}_x^*(\rho_\Lambda)(\log \rho_\Lambda - \log \sigma_\Lambda)]}{2D_x(\rho_\Lambda || \sigma_\Lambda)},$$

where σ_Λ is the fixed point of the evolution, and $D_x(\rho_\Lambda || \sigma_\Lambda)$ is the conditional relative entropy.

LEMMA

$$\alpha_\Lambda(\mathcal{L}_x^*) \geq \frac{1}{2}.$$

LOG-SOBOLEV CONSTANT

CONDITIONAL LOG-SOBOLEV CONSTANT

For $x \in \Lambda$, we define the **conditional log-Sobolev constant** of \mathcal{L}_Λ^* in x by

$$\alpha_\Lambda(\mathcal{L}_x^*) := \inf_{\rho_\Lambda \in \mathcal{S}_\Lambda} \frac{-\operatorname{tr}[\mathcal{L}_x^*(\rho_\Lambda)(\log \rho_\Lambda - \log \sigma_\Lambda)]}{2D_x(\rho_\Lambda || \sigma_\Lambda)},$$

where σ_Λ is the fixed point of the evolution, and $D_x(\rho_\Lambda || \sigma_\Lambda)$ is the conditional relative entropy.

LEMMA

$$\alpha_\Lambda(\mathcal{L}_x^*) \geq \frac{1}{2}.$$

POSITIVE LOG-SOBOLEV CONSTANT

$$\alpha(\mathcal{L}_\Lambda^*) \geq \frac{1}{2}.$$

LOG-SOBOLEV CONSTANT

CONDITIONAL LOG-SOBOLEV CONSTANT

For $x \in \Lambda$, we define the **conditional log-Sobolev constant** of \mathcal{L}_Λ^* in x by

$$\alpha_\Lambda(\mathcal{L}_x^*) := \inf_{\rho_\Lambda \in \mathcal{S}_\Lambda} \frac{-\operatorname{tr}[\mathcal{L}_x^*(\rho_\Lambda)(\log \rho_\Lambda - \log \sigma_\Lambda)]}{2D_x(\rho_\Lambda || \sigma_\Lambda)},$$

where σ_Λ is the fixed point of the evolution, and $D_x(\rho_\Lambda || \sigma_\Lambda)$ is the conditional relative entropy.

LEMMA

$$\alpha_\Lambda(\mathcal{L}_x^*) \geq \frac{1}{2}.$$

POSITIVE LOG-SOBOLEV CONSTANT

$$\alpha(\mathcal{L}_\Lambda^*) \geq \frac{1}{2}.$$

OPEN PROBLEMS

PROBLEM 1

Can we use any of the quasi-factorization results to prove log-Sobolev constants in a more general setting?

(**Kastoryano-Brandao, '15**) The heat-bath dynamics, with σ_Λ the Gibbs state of a commuting Hamiltonian, has positive spectral gap. \Rightarrow Log-Sobolev constant?

PROBLEM 2

Is there a better definition for conditional relative entropy?

PROBLEM 3

When do $D_A(\rho_{AB}||\sigma_{AB})$ and $D_A^E(\rho_{AB}||\sigma_{AB})$ coincide?

THANK YOU!

OPEN PROBLEMS

PROBLEM 1

Can we use any of the quasi-factorization results to prove log-Sobolev constants in a more general setting?

(**Kastoryano-Brandao, '15**) The heat-bath dynamics, with σ_Λ the Gibbs state of a commuting Hamiltonian, has positive spectral gap. \Rightarrow Log-Sobolev constant?

PROBLEM 2

Is there a better definition for conditional relative entropy?

PROBLEM 3

When do $D_A(\rho_{AB}||\sigma_{AB})$ and $D_A^E(\rho_{AB}||\sigma_{AB})$ coincide?

THANK YOU!