

# QUASI-FACTORIZATION OF THE QUANTUM RELATIVE ENTROPY

ÁNGELA CAPEL CUEVAS  
(ICMAT and U. Autónoma de Madrid)

Workshop on Quantum Functional Inequalities, Toulouse

Joint work with Angelo Lucia and David Pérez-García

arXiv: 1705.03521 and 1804.09525

12 June 2018

## 1 INTRODUCTION

## 2 CLASSICAL CASE

## 3 CONDITIONAL RELATIVE ENTROPY

- CONDITIONAL RELATIVE ENTROPY
- QUASI-FACTORIZATION FOR THE CONDITIONAL RELATIVE ENTROPY

## 4 CONDITIONAL RELATIVE ENTROPY BY EXPECTATIONS

- CONDITIONAL RELATIVE ENTROPY BY EXPECTATIONS
- QUASI-FACTORIZATION FOR THE CRE BY EXPECTATIONS

## 5 QUANTUM SPIN LATTICES

# 1. INTRODUCTION

- $\mathcal{H}_{AB} = \mathcal{H}_A \otimes \mathcal{H}_B$  (or  $\mathcal{H}_{ABC} = \mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$ ).
- $\mathcal{B}_\Lambda := \mathcal{B}(\mathcal{H}_\Lambda)$ , set of bounded linear operators.
- $\mathcal{A}_\Lambda \subseteq \mathcal{B}_\Lambda$ , set of Hermitian operators.
- $\mathcal{S}_\Lambda := \{f \in \mathcal{A}_\Lambda : f \geq 0 \text{ and } \text{tr}[f] = 1\}$ .
- $f \in \mathcal{B}_\Lambda$  has support on  $A \subseteq \Lambda$  if  $f = f_A \otimes \mathbb{1}_B$  for certain  $f_A \in \mathcal{B}_A$ .
- Modified partial trace:  $\text{tr}_A : f \mapsto \text{tr}_A[f] \otimes \mathbb{1}_A$ , where  $\text{tr}_A[f]$  has support in  $B$ .
- We denote by  $f_B$  the observable  $\text{tr}_A[f]$  with support in  $B$ .

## MIXING TIME

We define the **mixing time** of  $\{\mathcal{T}_t^*\}$  by

$$\tau(\varepsilon) = \min \left\{ t > 0 : \sup_{\rho \in \mathcal{S}_\Lambda} \|\mathcal{T}_t^*(\rho) - \mathcal{T}_\infty^*(\rho)\|_1 \leq \varepsilon \right\}.$$

## RAPID MIXING

We say that  $\mathcal{L}^*$  satisfies **rapid mixing** if

$$\sup_{\rho \in \mathcal{S}_\Lambda} \|\rho_t - \sigma\|_1 \leq \text{poly}(|\Lambda|) e^{-\gamma t}.$$

## PROBLEM

Find bounds for the mixing time!

## LOG-SOBOLEV CONSTANT

Let  $\mathcal{L} : \mathcal{B}_\Lambda \rightarrow \mathcal{B}_\Lambda$  be a primitive reversible Lindbladian with stationary state  $\sigma_\Lambda$ . We define the **log-Sobolev constant** (MLSI constant) of  $\mathcal{L}_\Lambda^*$  by

$$\alpha(\mathcal{L}_\Lambda^*) := \inf_{\rho_\Lambda \in \mathcal{S}_\Lambda} \frac{-\operatorname{tr}[\mathcal{L}_\Lambda^*(\rho_\Lambda)(\log \rho_\Lambda - \log \sigma_\Lambda)]}{2D(\rho_\Lambda || \sigma_\Lambda)}$$

We have:

$$\|\rho_t - \sigma_\Lambda\|_1 \leq \sqrt{2D(\rho_\Lambda || \sigma_\Lambda)} e^{-\alpha(\mathcal{L}_\Lambda^*)t} \leq \sqrt{2 \log(1/\sigma_{\min})} e^{-\alpha(\mathcal{L}_\Lambda^*)t}. \quad (1)$$

## RESULT

If  $\alpha(\mathcal{L}_\Lambda^*) > 0$ ,

$$\|\rho_t - \sigma_\Lambda\|_1 \leq \sqrt{2 \log(1/\sigma_{\min})} e^{-\alpha(\mathcal{L}_\Lambda^*)t}.$$

Log-Sobolev inequality  $\Rightarrow$  Rapid mixing.

## PROBLEM

Find positive log-Sobolev constants!

## 2. CLASSICAL CASE



## CLASSICAL ENTROPY AND CONDITIONAL ENTROPY

Consider a probability space  $(\Omega, \mathcal{F}, \mu)$  and define, for every  $f > 0$ , the **entropy** of  $f$  by

$$\text{Ent}_\mu(f) = \mu(f \log f) - \mu(f) \log \mu(f).$$

Given a  $\sigma$ -algebra  $\mathcal{G} \subseteq \mathcal{F}$ , we define the **conditional entropy** of  $f$  in  $\mathcal{G}$  by

$$\text{Ent}_\mu(f | \mathcal{G}) = \mu(f \log f | \mathcal{G}) - \mu(f | \mathcal{G}) \log \mu(f | \mathcal{G}).$$

With these definitions, the following lemma is proven:

LEMMA, Dai Pra et al. '02

Let  $(\Omega, \mathcal{F}, \mu)$  be a probability space, and  $\mathcal{F}_1, \mathcal{F}_2$  sub- $\sigma$ -algebras of  $\mathcal{F}$ . Suppose that there exists a probability measure  $\bar{\mu}$  that makes  $\mathcal{F}_1$  and  $\mathcal{F}_2$  independent,  $\mu \ll \bar{\mu}$  and  $\mu | \mathcal{F}_i = \bar{\mu} | \mathcal{F}_i$  for  $i = 1, 2$ . Then, for every  $f \geq 0$  such that  $f \log f \in L^1(\mu)$  and  $\mu(f) = 1$ ,

$$\text{Ent}_{\mu}(f) \leq \frac{1}{1 - 4\|h - 1\|_{\infty}} \mu [\text{Ent}_{\mu}(f | \mathcal{F}_1) + \text{Ent}_{\mu}(f | \mathcal{F}_2)],$$

where  $h = \frac{d\mu}{d\bar{\mu}}$ .

## PROBLEM

Let  $\mathcal{H}_{ABC} = \mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$  and  $\rho_{ABC}, \sigma_{ABC} \in \mathcal{S}_{ABC}$ . Can we prove something like

$$D(\rho_{ABC} || \sigma_{ABC}) \leq \xi(\sigma_{AB}) [D_{AB}(\rho_{ABC} || \sigma_{ABC}) + D_{BC}(\rho_{ABC} || \sigma_{ABC})] ?$$

Yes! (We will see how later)

### 3. CONDITIONAL RELATIVE ENTROPY

## QUANTUM RELATIVE ENTROPY

Let  $f, g \in \mathcal{A}_\Lambda$ ,  $f$  verifying  $\text{tr}[f] \neq 0$ . The **quantum relative entropy** of  $f$  and  $g$  is defined by:

$$D(f||g) = \frac{1}{\text{tr}[f]} \text{tr} [f(\log f - \log g)]. \quad (2)$$

## REMARK

In this talk, we only consider density matrices (with trace 1). In this case, the **quantum relative entropy** is given by:

$$D(\rho||\sigma) = \text{tr} [\rho(\log \rho - \log \sigma)]. \quad (3)$$

## PROPERTIES OF THE RELATIVE ENTROPY

Let  $\mathcal{H}_{AB} = \mathcal{H}_A \otimes \mathcal{H}_B$  and  $\rho_{AB}, \sigma_{AB} \in \mathcal{S}_{AB}$ . The following properties hold:

- 1 **Continuity.**  $\rho_{AB} \mapsto D(\rho_{AB}||\sigma_{AB})$  is continuous.
- 2 **Additivity.**  $D(\rho_A \otimes \rho_B || \sigma_A \otimes \sigma_B) = D(\rho_A || \sigma_A) + D(\rho_B || \sigma_B)$ .
- 3 **Superadditivity.**  

$$D(\rho_{AB} || \sigma_A \otimes \sigma_B) \geq D(\rho_A || \sigma_A) + D(\rho_B || \sigma_B).$$
- 4 **Monotonicity.**  $D(\rho_{AB} || \sigma_{AB}) \geq D(T(\rho_{AB}) || T(\sigma_{AB}))$  for every quantum channel  $T$ .

## CHARACTERIZATION OF THE RELATIVE ENTROPY, Wilming et al. '17

If  $f : \mathcal{S}_{AB} \times \mathcal{S}_{AB} \rightarrow \mathbb{R}_0^+$  satisfies 1 – 4, then  $f$  is the relative entropy.

# CONDITIONAL RELATIVE ENTROPY

## CONDITIONAL RELATIVE ENTROPY

Let  $\mathcal{H}_{AB} = \mathcal{H}_A \otimes \mathcal{H}_B$ . We define a **conditional relative entropy** in  $A$  as a function

$$D_A(\cdot || \cdot) : \mathcal{S}_{AB} \times \mathcal{S}_{AB} \rightarrow \mathbb{R}_0^+$$

verifying the following properties for every  $\rho_{AB}, \sigma_{AB} \in \mathcal{S}_{AB}$ :

- 1 **Continuity:** The map  $\rho_{AB} \mapsto D_A(\rho_{AB} || \sigma_{AB})$  is continuous.
- 2 **Non-negativity:**  $D_A(\rho_{AB} || \sigma_{AB}) \geq 0$  and  
 (2.1)  $D_A(\rho_{AB} || \sigma_{AB}) = 0$  if, and only if,  $\rho_{AB} = \mathbb{E}_A^*(\rho_{AB})$ .
- 3 **Semi-superadditivity:**  $D_A(\rho_{AB} || \sigma_A \otimes \sigma_B) \geq D(\rho_A || \sigma_A)$  and  
 (3.1) **Semi-additivity:** if  $\rho_{AB} = \rho_A \otimes \rho_B$ ,  
 $D_A(\rho_A \otimes \rho_B || \sigma_A \otimes \sigma_B) = D(\rho_A || \sigma_A)$ .
- 4 **Semi-monotonicity:** For every quantum channel  $\mathcal{T}$ ,  
 $D_A(\mathcal{T}(\rho_{AB}) || \mathcal{T}(\sigma_{AB})) + D_B((\text{tr}_A \circ \mathcal{T})(\rho_{AB}) || (\text{tr}_A \circ \mathcal{T})(\sigma_{AB}))$   
 $\leq D_A(\rho_{AB} || \sigma_{AB}) + D_B(\text{tr}_A(\rho_{AB}) || \text{tr}_A(\sigma_{AB})).$

## REMARK

Consider for every  $\rho_{AB}, \sigma_{AB} \in \mathcal{S}_{AB}$

$$D_{A,B}^+(\rho_{AB}||\sigma_{AB}) = D_A(\rho_{AB}||\sigma_{AB}) + D_B(\rho_{AB}||\sigma_{AB}).$$

Then,  $D_{A,B}^+$  verifies the following properties:

① **Continuity:**  $\rho_{AB} \mapsto D_{A,B}^+(\rho_{AB}||\sigma_{AB})$  is continuous.

② **Additivity:**

$$D_{A,B}^+(\rho_A \otimes \rho_B || \sigma_A \otimes \sigma_B) = D(\rho_A || \sigma_A) + D(\rho_B || \sigma_B).$$

③ **Superadditivity:**

$$D_{A,B}^+(\rho_{AB} || \sigma_A \otimes \sigma_B) \geq D(\rho_A || \sigma_A) + D(\rho_B || \sigma_B).$$

However, it does not satisfy the property of monotonicity.




## AXIOMATIC CHARACTERIZATION OF THE CONDITIONAL RELATIVE ENTROPY

The only possible conditional relative entropy is given by:

$$D_A(\rho_{AB}||\sigma_{AB}) = D(\rho_{AB}||\sigma_{AB}) - D(\rho_B||\sigma_B)$$

for every  $\rho_{AB}, \sigma_{AB} \in \mathcal{S}_{AB}$ .



**Figura:** Choice of indices in  $\mathcal{H}_{ABC} = \mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$ .

Result of **quasi-factorization** of the relative entropy, for every  $\rho_{ABC}, \sigma_{ABC} \in \mathcal{S}_{ABC}$ :

$$(1 - \xi(\sigma_{ABC}))D(\rho_{ABC}||\sigma_{ABC}) \leq D_{AB}(\rho_{ABC}||\sigma_{ABC}) + D_{BC}(\rho_{ABC}||\sigma_{ABC}),$$

where  $\xi(\sigma_{ABC})$  depends only on  $\sigma_{ABC}$  and measures how far  $\sigma_{AC}$  is from  $\sigma_A \otimes \sigma_C$ .

## QUASI-FACTORIZATION FOR THE CRE

Let  $\mathcal{H}_{ABC} = \mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$  and  $\rho_{ABC}, \sigma_{ABC} \in \mathcal{S}_{ABC}$ . Then, the following inequality holds

$$(1 - 2\|H(\sigma_{AC})\|_\infty)D(\rho_{ABC}||\sigma_{ABC}) \leq D_{AB}(\rho_{ABC}||\sigma_{ABC}) + D_{BC}(\rho_{ABC}||\sigma_{ABC}),$$

where

$$H(\sigma_{AC}) = \sigma_A^{-1/2} \otimes \sigma_C^{-1/2} \sigma_{AC} \sigma_A^{-1/2} \otimes \sigma_C^{-1/2} - \mathbb{1}_{AC}.$$

Note that  $H(\sigma_{AC}) = 0$  if  $\sigma_{AC}$  is a tensor product between  $A$  and  $C$ .

$$\begin{aligned}
 (1 - 2\|H(\sigma_{AC})\|_\infty)D(\rho_{ABC}||\sigma_{ABC}) &\leq \\
 D_{AB}(\rho_{ABC}||\sigma_{ABC}) + D_{BC}(\rho_{ABC}||\sigma_{ABC}) &= \\
 = 2D(\rho_{ABC}||\sigma_{ABC}) - D(\rho_C||\sigma_C) - D(\rho_A||\sigma_A). &
 \end{aligned}$$

$$\Leftrightarrow$$

$$\begin{aligned}
 (1 + 2\|H(\sigma_{AC})\|_\infty)D(\rho_{ABC}||\sigma_{ABC}) &\geq \\
 D(\rho_A||\sigma_A) + D(\rho_C||\sigma_C). &
 \end{aligned}$$

$$\Leftrightarrow$$

$$\begin{aligned}
 (1 + 2\|H(\sigma_{AC})\|_\infty)D(\rho_{AC}||\sigma_{AC}) &\geq \\
 D(\rho_A||\sigma_A) + D(\rho_C||\sigma_C). &
 \end{aligned}$$

Recall:

- **Superadditivity.**

$$D(\rho_{AB}||\sigma_A \otimes \sigma_B) \geq D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B).$$

Due to:

- **Monotonicity.**  $D(\rho_{AB}||\sigma_{AB}) \geq D(T(\rho_{AB})||T(\sigma_{AB}))$  for every quantum channel  $T$ .

we have

$$2D(\rho_{AB}||\sigma_{AB}) \geq D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B)$$

Our result:

$$\boxed{(1 + 2\|H(\sigma_{AB})\|_\infty)D(\rho_{AB}||\sigma_{AB}) \geq D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B)}.$$

## QUASI-FACTORIZATION FOR THE CRE

Let  $\mathcal{H}_{ABC} = \mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$  and  $\rho_{ABC}, \sigma_{ABC} \in \mathcal{S}_{ABC}$ . Then, the following inequality holds

$$(1 - 2\|H(\sigma_{AC})\|_\infty)D(\rho_{ABC}||\sigma_{ABC}) \leq D_{AB}(\rho_{ABC}||\sigma_{ABC}) + D_{BC}(\rho_{ABC}||\sigma_{ABC}),$$

where

$$H(\sigma_{AC}) = \sigma_A^{-1/2} \otimes \sigma_C^{-1/2} \sigma_{AC} \sigma_A^{-1/2} \otimes \sigma_C^{-1/2} - \mathbb{1}_{AC}.$$

Note that  $H(\sigma_{AC}) = 0$  if  $\sigma_{AC}$  is a tensor product between  $A$  and  $C$ .

## STEP 1

$$D(\rho_{AB}||\sigma_{AB}) \geq D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B) - \log \operatorname{tr} M, \quad (4)$$

where  $M = \exp [\log \sigma_{AB} - \log \sigma_A \otimes \sigma_B + \log \rho_A \otimes \rho_B]$ .

It holds that:

$$\begin{aligned} & D(\rho_{AB}||\sigma_{AB}) - [D(\rho_A||\sigma_A) + D(\rho_B||\sigma_B)] = \\ & = \operatorname{tr} \left[ \rho_{AB} \left( \log \rho_{AB} - \underbrace{(\log \sigma_{AB} - \log \sigma_A \otimes \sigma_B + \log \rho_A \otimes \rho_B)}_{\log M} \right) \right] \\ & = D(\rho_{AB}||M) \geq -\log \operatorname{tr} M. \end{aligned}$$

## STEP 2

$$\log \operatorname{tr} M \leq \operatorname{tr}[L(\sigma_{AB})(\rho_A - \sigma_A) \otimes (\rho_B - \sigma_B)], \quad (5)$$

where

$$L(\sigma_{AB}) = \mathcal{T}_{\sigma_A \otimes \sigma_B}(\sigma_{AB}) - \mathbb{1}_{AB}.$$



## THEOREM (LIEB)

Let  $g$  a positive operator, and define

$$\mathcal{T}_g(f) = \int_0^\infty dt (g+t)^{-1} f (g+t)^{-1}.$$

$\mathcal{T}_g$  is positive-semidefinite if  $g$  is. We have that

$$\mathrm{tr}[\exp(-f + g + h)] \leq \mathrm{tr}[e^h \mathcal{T}_{ef}(e^g)].$$

We apply Lieb's theorem to the previous equation :

$$\begin{aligned} \mathrm{tr} M &\leq \mathrm{tr}[\rho_A \otimes \rho_B \mathcal{T}_{\sigma_A \otimes \sigma_B}(\sigma_{AB})] \\ &= \mathrm{tr} \left[ \rho_A \otimes \rho_B \underbrace{(\mathcal{T}_{\sigma_A \otimes \sigma_B}(\sigma_{AB}) - \mathbb{1}_{AB})}_{L(\sigma_{AB})} \right] + \underbrace{\mathrm{tr}[\rho_A \otimes \rho_B]}_1. \end{aligned}$$

By using the fact  $\log(x) \leq x - 1$ , we conclude

$$\log \mathrm{tr} M \leq \mathrm{tr} M - 1 \leq \mathrm{tr}[L(\sigma_{AB}) \rho_A \otimes \rho_B].$$

## LEMMA (SUTTER ET AL.)

For  $f \in \mathcal{S}_{AB}$  and  $g \in \mathcal{A}_{AB}$  the following holds:

$$\mathcal{T}_g(f) = \int_{-\infty}^{\infty} dt \beta_0(t) g^{\frac{-1-it}{2}} f g^{\frac{-1+it}{2}},$$

with

$$\beta_0(t) = \frac{\pi}{2} (\cosh(\pi t) + 1)^{-1}.$$

## LEMMA

For every operator  $O_A \in \mathcal{B}_A$  and  $O_B \in \mathcal{B}_B$  the following holds:

$$\text{tr}[L(\sigma_{AB}) \sigma_A \otimes O_B] = \text{tr}[L(\sigma_{AB}) O_A \otimes \sigma_B] = 0.$$

## STEP 3

$$\mathrm{tr}[L(\sigma_{AB})(\rho_A - \sigma_A) \otimes (\rho_B - \sigma_B)] \leq 2\|L(\sigma_{AB})\|_\infty D(\rho_{AB}||\sigma_{AB}). \quad (6)$$

In virtue of Hölder's inequality and tensorization of Schatten norms,

$$\begin{aligned} \mathrm{tr}[L(\sigma_{AB})(\rho_A - \sigma_A) \otimes (\rho_B - \sigma_B)] &\leq \\ &\|L(\sigma_{AB})\|_\infty \|(\rho_A - \sigma_A) \otimes (\rho_B - \sigma_B)\|_1 \\ &= \|L(\sigma_{AB})\|_\infty \|\rho_A - \sigma_A\|_1 \|\rho_B - \sigma_B\|_1. \end{aligned}$$

### THEOREM (PINSKER)

For  $\rho_{AB}$  and  $\sigma_{AB}$  density matrices, it holds that

$$\|\rho_{AB} - \sigma_{AB}\|_1^2 \leq 2D(\rho_{AB}||\sigma_{AB}).$$

Using Pinsker's theorem and the data-processing inequality, we can conclude:

$$\text{tr}[L(\sigma_{AB})(\rho_A - \sigma_A) \otimes (\rho_B - \sigma_B)] \leq 2\|L(\sigma_{AB})\|_\infty D(\rho_{AB}||\sigma_{AB}).$$

## Step 4

$$\|L(\sigma_{AB})\|_{\infty} \leq \left\| \sigma_A^{-1/2} \otimes \sigma_B^{-1/2} \sigma_{AB} \sigma_A^{-1/2} \otimes \sigma_B^{-1/2} - \mathbb{1}_{AB} \right\|_{\infty}. \quad (7)$$

## 4. CONDITIONAL RELATIVE ENTROPY BY EXPECTATIONS

## WEAK CONDITIONAL RELATIVE ENTROPY

Let  $\mathcal{H}_{AB} = \mathcal{H}_A \otimes \mathcal{H}_B$ . We define a **weak conditional relative entropy** in  $A$  as a function

$$D_A(\cdot || \cdot) : \mathcal{S}_{AB} \times \mathcal{S}_{AB} \rightarrow \mathbb{R}_0^+$$

verifying the following properties for every  $\rho_{AB}, \sigma_{AB} \in \mathcal{S}_{AB}$ :

- 1 **Continuity:** The map  $\rho_{AB} \mapsto D_A(\rho_{AB} || \sigma_{AB})$  is continuous.
- 2 **Non-negativity:**  $D_A(\rho_{AB} || \sigma_{AB}) \geq 0$  and  
 (2.1)  $D_A(\rho_{AB} || \sigma_{AB}) = 0$  if, and only if,  $\rho_{AB} = \mathbb{E}_A^*(\rho_{AB})$ .
- 3 **Semi-superadditivity:**  $D_A(\rho_{AB} || \sigma_A \otimes \sigma_B) \geq D(\rho_A || \sigma_A)$  and  
 (3.1) **Semi-additivity:** if  $\rho_{AB} = \rho_A \otimes \rho_B$ ,  
 $D_A(\rho_A \otimes \rho_B || \sigma_A \otimes \sigma_B) = D(\rho_A || \sigma_A)$ .

## MINIMAL CONDITIONAL EXPECTATION

Let  $\mathcal{H}_{AB} = \mathcal{H}_A \otimes \mathcal{H}_B$  and  $\sigma_{AB} \in \mathcal{S}_{AB}$ ,  $f_{AB} \in \mathcal{A}_{AB}$ . We define the **minimal conditional expectation** of  $\sigma_{AB}$  on  $A$  by

$$\mathbb{E}_A^\sigma(f_{AB}) := \text{tr}_A[\eta_A^\sigma f_{AB} \eta_A^{\sigma\dagger}], \quad (8)$$

where  $\eta_A^\sigma := (\text{tr}_A[\sigma_{AB}])^{-1/2} \sigma_{AB}^{1/2}$ .

For  $\rho_{AB} \in \mathcal{S}_{AB}$ ,  $(\mathbb{E}_A^\sigma)^*$  (hereafter denoted by  $\mathbb{E}_A^*$ ) is given by

$$\mathbb{E}_A^*(\rho_{AB}) := \sigma_{AB}^{1/2} \sigma_B^{-1/2} \rho_B \sigma_B^{-1/2} \sigma_{AB}^{1/2}. \quad (9)$$

It coincides with the Petz recovery map for the partial trace.



# CONDITIONAL RELATIVE ENTROPY BY EXPECTATIONS

## CONDITIONAL RELATIVE ENTROPY BY EXPECTATIONS

Let  $\mathcal{H}_{AB} = \mathcal{H}_A \otimes \mathcal{H}_B$  and  $\rho_{AB}, \sigma_{AB} \in S_{AB}$ . Let  $\mathbb{E}_A^*$  be defined as above. We define the **conditional relative entropy by expectations** of  $\rho_{AB}$  and  $\sigma_{AB}$  in  $A$  by:

$$D_A^E(\rho_{AB} || \sigma_{AB}) = D(\rho_{AB} || \mathbb{E}_A^*(\rho_{AB})).$$

## PROPERTY

$D_A^E(\rho_{AB} || \sigma_{AB})$  is a weak conditional relative entropy.

## PROBLEM

Under which conditions holds

$$D_A(\rho_{AB}||\sigma_{AB}) = D_A^E(\rho_{AB}||\sigma_{AB})?$$

## EXAMPLES

- ① If  $[\rho_B, \sigma_{AB}] = [\rho_B, \sigma_B] = [\sigma_B, \sigma_{AB}] = 0$ ,

$$D_A(\rho_{AB}||\sigma_{AB}) = D_A^E(\rho_{AB}||\sigma_{AB}).$$

- ② If  $\sigma = \sigma_A \otimes \sigma_B$ , then

$$D_A(\rho_{AB}||\sigma_{AB}) = D_A^E(\rho_{AB}||\sigma_{AB}).$$

- ③  $D_A(\rho_{AB}||\sigma_{AB}) = 0 \Leftrightarrow D_A^E(\rho_{AB}||\sigma_{AB}) = 0$ .

In general, it is an open question.

## RELATION WITH THE CLASSICAL CASE




Figura: Identification between classical and quantum quantities when the states considered are classical.

## QUASI-FACTORIZATION CRE BY EXPECTATIONS

Let  $\mathcal{H}_{AB} = \mathcal{H}_A \otimes \mathcal{H}_B$  and  $\rho_{AB}, \sigma_{AB} \in \mathcal{S}_{AB}$ . The following inequality holds

$$(1 - \xi(\sigma_{AB}))D(\rho_{AB}||\sigma_{AB}) \leq D_A^E(\rho_{AB}||\sigma_{AB}) + D_B^E(\rho_{AB}||\sigma_{AB}), \quad (10)$$

where

$$\xi(\sigma_{AB}) = 2(E_1(t) + E_2(t)),$$

and

$$E_1(t) = \int_{-\infty}^{+\infty} dt \beta_0(t) \left\| \sigma_B^{\frac{-1+it}{2}} \sigma_{AB}^{\frac{1-it}{2}} \sigma_A^{\frac{-1+it}{2}} - \mathbb{1}_{AB} \right\|_{\infty} \left\| \sigma_A^{-1/2} \sigma_{AB}^{\frac{1+it}{2}} \sigma_B^{-1/2} \right\|_{\infty},$$

$$E_2(t) = \int_{-\infty}^{+\infty} dt \beta_0(t) \left\| \sigma_B^{\frac{-1-it}{2}} \sigma_{AB}^{\frac{1+it}{2}} \sigma_A^{\frac{-1-it}{2}} - \mathbb{1}_{AB} \right\|_{\infty}.$$

Note that  $\xi(\sigma_{AB}) = 0$  if  $\sigma_{AB}$  is a tensor product between  $A$  and  $B$ .

## 5. QUANTUM SPIN LATTICES

## QUANTUM SPIN LATTICES





Figura: A quantum spin lattice system.

- Lattice  $\Lambda \subset \mathbb{Z}^d$ .
- For every site  $x$ ,  $\mathcal{H}_x (= \mathbb{C}^D)$ .
- The global Hilbert space associated to  $\Lambda$  is  $\mathcal{H}_\Lambda = \bigotimes_{x \in \Lambda} \mathcal{H}_x$ .

## QUANTUM SPIN LATTICES



**Figura:** A quantum spin lattice system  $\Lambda$  and  $A, B \subseteq \Lambda$  such that  $A \cup B = \Lambda$ .

GENERAL QUASI-FACTORIZATION FOR  $\sigma$  A TENSOR PRODUCT

Let  $\mathcal{H}_\Lambda = \bigotimes_{x \in \Lambda} \mathcal{H}_x$  and  $\rho_\Lambda, \sigma_\Lambda \in \mathcal{S}_\Lambda$  such that  $\sigma_\Lambda = \bigotimes_{x \in \Lambda} \sigma_x$ . The following inequality holds:

$$D(\rho_\Lambda || \sigma_\Lambda) \leq \sum_{x \in \Lambda} D_x(\rho_\Lambda || \sigma_\Lambda). \quad (11)$$

Proof based on **strong subadditivity**.



The **heat-bath dynamics**, with product fixed point, has a positive log-Sobolev constant.

Consider the local Lindbladian

$$\mathcal{L}_x^* := \mathbb{E}_x^* - \mathbb{1}_\Lambda,$$

and the global Lindbladian

$$\mathcal{L}_\Lambda^* = \sum_{x \in \Lambda} \mathcal{L}_x^*.$$

Since

$$\mathbb{E}_x^*(\rho_\Lambda) = \sigma_\Lambda^{1/2} \sigma_{x^c}^{-1/2} \rho_{x^c} \sigma_{x^c}^{-1/2} \sigma_\Lambda^{1/2} = \sigma_x \otimes \rho_{x^c}$$

for every  $\rho_\Lambda \in \mathcal{S}_\Lambda$ , we have

$$\mathcal{L}_\Lambda^*(\rho_\Lambda) = \sum_{x \in \Lambda} (\sigma_x \otimes \rho_{x^c} - \rho_\Lambda).$$

## CONDITIONAL LOG-SOBOLEV CONSTANT

For  $A \subset \Lambda$ , we define the **conditional log-Sobolev constant** of  $\mathcal{L}_\Lambda^*$  in  $A$  by

$$\alpha_\Lambda(\mathcal{L}_A^*) := \inf_{\rho_\Lambda \in \mathcal{S}_\Lambda} \frac{-\operatorname{tr}[\mathcal{L}_A^*(\rho_\Lambda)(\log \rho_\Lambda - \log \sigma_\Lambda)]}{2D_A(\rho_\Lambda \parallel \sigma_\Lambda)},$$

where  $\sigma_\Lambda$  is the fixed point of the evolution, and  $D_A(\rho_\Lambda \parallel \sigma_\Lambda)$  is the conditional relative entropy.

## LEMMA

$$\alpha_\Lambda(\mathcal{L}_x^*) \geq \frac{1}{2}.$$

## POSITIVE LOG-SOBOLEV CONSTANT

$$\alpha(\mathcal{L}_\Lambda^*) \geq \frac{1}{2}.$$

$$\begin{aligned}
 D(\rho_\Lambda || \sigma_\Lambda) &\leq \sum_{x \in \Lambda} D_x(\rho_\Lambda || \sigma_\Lambda) \\
 &\leq \sum_{x \in \Lambda} \frac{-\operatorname{tr}[\mathcal{L}_x^*(\rho_\Lambda)(\log \rho_\Lambda - \log \sigma_\Lambda)]}{2\alpha_\Lambda(\mathcal{L}_x^*)} \\
 &\leq \frac{1}{2 \inf_{x \in \Lambda} \alpha_\Lambda(\mathcal{L}_x^*)} \sum_{x \in \Lambda} -\operatorname{tr}[\mathcal{L}_x^*(\rho_\Lambda)(\log \rho_\Lambda - \log \sigma_\Lambda)] \\
 &= \frac{1}{2 \inf_{x \in \Lambda} \alpha_\Lambda(\mathcal{L}_x^*)} (-\operatorname{tr}[\mathcal{L}_\Lambda^*(\rho_\Lambda)(\log \rho_\Lambda - \log \sigma_\Lambda)]) \\
 &\leq (-\operatorname{tr}[\mathcal{L}_\Lambda^*(\rho_\Lambda)(\log \rho_\Lambda - \log \sigma_\Lambda)]).
 \end{aligned}$$

FOR FURTHER KNOWLEDGE,  
ARXIV: 1705.03521 AND 1804.09525



thank you!